

Oral presentation

Specificity of commonly used cyclic nucleotides with PKA, PKG and Epac-implementing microcalorimetry to determine PDE activities

Joseph A Beavo¹, Johannes L Bos², Elke Butt^{*3}, Stein O Doskeland⁵, Hans-Gottfried Genieser⁴, Thomas R Hinds¹, Heiko Poppe^{1,3}, Holger Rehmann², Sergei D Rybalkin¹, Frank Schwede⁴ and Xiao-Bo Tang¹

Address: ¹Department of Pharmacology, University of Washington, Seattle, WA 98195, USA, ²Department of Physiological Chemistry and Centre for Biomedical Genetics, University Medical Center Utrecht, 3584 CG Utrecht, The Netherlands, ³Institute of Clinical Biochemistry, University of Wuerzburg, Wuerzburg, Germany, ⁴Biolog. Life Science Institute, Flughafendamm 9a, 28199 Bremen, Germany and ⁵Department of Anatomy and Cell Biology, University of Bergen, N-5009, Bergen, Norway

Email: Elke Butt* - butt@klin-biochem.uni-wuerzburg.de

* Corresponding author

from 3rd International Conference on cGMP Generators, Effectors and Therapeutic Implications
Dresden, Germany. 15–17 June 2007

Published: 25 July 2007

BMC Pharmacology 2007, 7(Suppl 1):S25 doi:10.1186/1471-2210-7-S1-S25

This abstract is available from: <http://www.biomedcentral.com/1471-2210/7/S1/S25>

© 2007 Beavo et al; licensee BioMed Central Ltd.

Background

Two main structural classes of cyclic nucleotide receptors exist in mammalian cells. The first class contains a conserved cyclic nucleotide binding (CNB) domain, found in the regulatory regions of PKG and PKA in cyclic nucleotide regulated ion channels and in Epac. The second class contains a GAF domain that can bind cyclic nucleotides but is structurally unrelated to CNB domains. GAF domains are present in some phosphodiesterases (PDEs) allosterically regulated by cyclic nucleotides. In addition, cGMP and cAMP are bound and hydrolysed by PDEs at their catalytic site.

Materials and methods

The important feature of this work was to measure 12 of the most commonly used cyclic nucleotide analogues on protein kinase A I and II (PKA I and II), protein kinase G I α , I β and II (PKG I α , I β and II), and the guanine nucleotide exchange factor Epac I. The study is extended to eight cyclic nucleotide hydrolysing phosphodiesterases (PDE IA, IB, IC, II, IV, V, VI and X), thereby introducing microcalorimetry to overcome radioactive labeling of the tested analogs and allowing analysis of the kinetic parameters K_m , K_i and V_{max} [1].

Results and discussion

We successfully used isothermal titration microcalorimetry (ITC) for analysing the kinetic parameters of eight PDEs by various cAMP- and cGMP-analogs. The measured data reflect very well the reported literature and the technique offers significant advantages over traditional microbiological methods. The measuring of both, k_{cat} and K_m , provide additional insight into protein active sites by determining competitive, non-competitive and linear-mixed inhibition of the PDEs by the cyclic nucleotides.

To our surprise, some accepted hydrolysis-stable analogs became degraded by the newer phosphodiesterases or show high inhibitory effects. As an example, we show side effects of an Epac specific activator on platelet signalling pathways.

The defined specificity profiles of the tested cyclic nucleotides are presented in clearly arranged tables and will help to interpret *in vivo* data in the light of the cyclic nucleotide receptor proteins present in a given cell.

An extended overview of the data will be accessible to the scientific community at: <http://www.cyclic-nucleotides.net> after publication.

References

1. Downey C, Kazmi R, Toh CH: **Early identification and prognostic implications in disseminated intravascular coagulation through transmittance waveform analysis.** *Thromb Haemost* 1998, **80**:65-69.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

